ISE 2211 Statistics for Engineers

Homework #9 - Chapter 13 ANOVA (Montgomery & Runger, 5ed)

Reading Assignment: Pages 513 - 521

Problems:

Notes:

- 1. Use the numbered textbook problems only as references to help add meaning to the data.
- 2. Do NOT solve the problems as stated in the textbook.
- 3. For each of the problems, use ANOVA techniques to determine whether or not there is a statistically significant difference between the various treatments ($\alpha = 0.05$).
- 4. In order to save you from the drudgery of tedious arithmetic, the SS_{Total} and $SS_{Treatment}$ are given. The F_{test} and $F_{Critical Value}$ are also provided so that you may check your work. Work through the necessary calculations to determine the F_{test} and $F_{Critical Value}$ and then draw your conclusions.

Problem	$\mathbf{SS}_{_{\mathrm{Total}}}$	SS _{Treatment}	F _{test}	F _{Critical Value}
13 - 3	636.96	475.76	14.76	2.87
13 - 4	11.28	3.65	3.59	3.68
13 - 5	643648	489740	12.73	3.49
13 - 6	651.7	260.9	4.01	3.89
13 - 7	1303.7	1060.5	16.35	3.06
13 - 8	1265.6	1133.4	30.85	2.77
	Problem 13 - 3 13 - 4 13 - 5 13 - 6 13 - 7 13 - 8	Problem SS_{Total} 13 - 3636.9613 - 411.2813 - 564364813 - 6651.713 - 71303.713 - 81265.6	Problem SS_{Total} $SS_{Treatment}$ 13 - 3636.96475.7613 - 411.283.6513 - 564364848974013 - 6651.7260.913 - 71303.71060.513 - 81265.61133.4	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$

Conclusions:

13-3	Reject	Conclude, at least one sample significantly different.
13-4	Fail to Reject	Insufficient evidence to say at least one sample significantly different.
13-5	Reject	Conclude, at least one sample significantly different.
13-6	Reject	Conclude, at least one sample significantly different.
13-7	Reject	Conclude, at least one sample significantly different.
13-8	Reject	Conclude, at least one sample significantly different.